

One-Particle Excitation of the Two-Dimensional Hubbard Model*

Ken YOKOYAMA** and Hidetoshi FUKUYAMA

Department of Physics, Tokyo University, Tokyo 113

(Received January 21, 1997)

The real part of the self-energy of interacting two-dimensional electrons has been calculated in the t -matrix approximation. It is shown that the forward scattering results in an anomalous term leading to the vanishing renormalization factor of the one-particle Green function, which is a non-perturbative effect of the interaction U . The present result is a microscopic demonstration of the claim by Anderson based on the conventional many-body theory. The effect of the damping of the interacting electrons, which has been ignored in reaching above conclusion, has been briefly discussed.

KEYWORDS: Hubbard model, two dimensions, t -matrix approximation, forward scattering

The nature of the low-energy excitation of the two-dimensional system is of great interest recently. The Fermi-liquid picture was considered to be valid from diagrammatic studies,¹⁾ while it was suggested by Anderson that the anomalous behavior of the forward scattering phase shift leads to the renormalization factor $Z = 0$, i.e., the breakdown of the Fermi-liquid.^{2,3)} This remarkable suggestion has attracted much interest,^{4,5)} and several calculations of the self-energy of the two-dimensional Hubbard model have been carried out based on the t -matrix approximation,^{6,7,8,9)} in which the self-energy is approximated by the summation of ladder diagrams of the particle-particle process. In these calculations, however, only the imaginary part of the self-energy has been considered, and the real part has not been studied in detail.

In this paper, we will calculate explicitly the real part of the self-energy of the two-dimensional Hubbard model by the t -matrix approximation, and show that the singularity of the t -matrix in the forward scattering region gives rise to an anomalous term to the real part of the self-energy, which leads to the renormalization factor $Z = 0$. This result is in accordance with the claim by Anderson.

* To appear in J. Phys. Soc. Jpn. **66** No. 3 (1997).

** E-mail: yokoyama@watson.phys.s.u-tokyo.ac.jp

We consider the asymptotic behavior of the self-energy $\Sigma(\mathbf{k}, \epsilon + i\delta)$ in the limit of $|\mathbf{k} - \mathbf{k}_F| \ll k_F$, $|\epsilon| \ll \epsilon_F$, $\mathbf{k}_F$ and ϵ_F being the Fermi momentum and the Fermi energy. In general, the shape of the Fermi surface near the point $\mathbf{k}$ in the momentum space can be approximated as a parabolic curve, as is schematically shown in Fig. 1(a) and 1(b). Linearizing the energy dispersion in the normal direction of the Fermi surface, we assume the following energy dispersion.¹⁰⁾

$$\xi_k \equiv \epsilon_k - \mu \equiv v_0 k_x + \frac{A_0}{2} k_y^2, \quad (1)$$

where we take the origin of momentum at the point nearest to $\mathbf{k}$ on the Fermi surface, as is shown in Fig. 1(b), μ , v_0 and A_0 are the chemical potential, the Fermi-velocity and a constant, respectively. Here, the momentum dependence of the velocity is neglected. We calculate the

Fig. 1. The local shape of (a) the Fermi surface can be approximated by (b) a parabolic curve. (c) The energy ϵ dependence of the self-energy in the vicinity of $v\mathbf{k}$ is focused on.

self-energy $\Sigma(\mathbf{k}, \epsilon + i\delta)$ at the point $\mathbf{k} = (k, 0)$. In the Hubbard model, the t -matrix is given by

$$T(\mathbf{q}, x + i\delta) = \frac{-U}{1 + UK(\mathbf{q}, x + i\delta)} \quad (2)$$

$$K(\mathbf{q}, x + i\delta) = \sum_k \frac{\text{sgn} \xi_{k+q/2}}{\xi_{k+q/2} + \xi_{-k+q/2} - x - i\delta}, \quad (3)$$

where $K(\mathbf{q}, x + i\delta)$ is the particle-particle correlation function. Noting that $K(\mathbf{q}, 0)$ as a function of $\mathbf{q}$ is regular in the forward scattering region, $\mathbf{q} \simeq 2\mathbf{k}_F$, we introduce $K_0 \equiv \lim_{q \rightarrow 2k_F} [\lim_{x \rightarrow 0} K(\mathbf{q}, x + i\delta)]$, which has a contribution from the high-energy region, and reflects the whole band structure. In contrast to this, q - and x -dependences of $K(\mathbf{q}, x + i\delta) - K_0$ in the region of $|\mathbf{q} - 2\mathbf{k}_F| \ll k_F$, $|x| \ll \epsilon_F$ reflect details of a scattering process near the Fermi-energy. Hence, we write the t -matrix as,

$$T(\mathbf{q}, x + i\delta) = \frac{-U_{\text{eff}}}{1 + U_{\text{eff}} [K(\mathbf{q}, x + i\delta) - K_0]} \quad (4)$$

$$U_{\text{eff}} \equiv \frac{U}{1 + UK_0}. \quad (5)$$

We will focus only on the contribution of the t -matrix to $\Sigma(\mathbf{k}, \epsilon + i\delta)$ from the forward scattering region, i.e., the total momentum of the particle-particle ladder is nearly $2\mathbf{k}$. In the following calculations, we introduce momentum cut-offs k_c and k'_c for x and y components of the momentum in the energy dispersion, and momentum integrations for intermediate states of a scattering process are carried out within these cut-offs.

In the t -matrix approximation, the expressions for the real and imaginary parts of the self-energy are given by

$$\begin{aligned} \text{Re}\Sigma(\mathbf{k}, \epsilon + i\delta) &= \int \frac{d^2q}{(2\pi)^2} \int_{\epsilon}^{\infty} \frac{dx}{\pi} \text{Re}T(\mathbf{q}, x + i\delta) \text{Im}G_0(\mathbf{q} - \mathbf{k}, x - \epsilon - i\delta) \\ &\quad - \int \frac{d^2q}{(2\pi)^2} \int_0^{\infty} \frac{dx}{\pi} \text{Im}T(\mathbf{q}, x + i\delta) \text{Re}G_0(\mathbf{q} - \mathbf{k}, x - \epsilon - i\delta) \end{aligned} \quad (6)$$

$$\text{Im}\Sigma(\mathbf{k}, \epsilon + i\delta) = - \int \frac{d^2q}{(2\pi)^2} \int_0^{\epsilon} \frac{dx}{\pi} \text{Im}T(\mathbf{q}, x + i\delta) \text{Im}G_0(\mathbf{q} - \mathbf{k}, x - \epsilon - i\delta). \quad (7)$$

We will evaluate the asymptotic forms of $\text{Re}\Sigma(\mathbf{k}, \epsilon + i\delta)$ and $\text{Im}\Sigma(\mathbf{k}, \epsilon + i\delta)$ under the condition of $|\epsilon|, |vk| \ll \epsilon_c$, where ϵ_c is defined as $\text{Min}(v_0 k_c, A_0 k'_c{}^2)$. Dividing x integration in $[\epsilon, \infty]$ in the first term of eq. (6) into $[0, \epsilon]$ and $[0, \infty]$, we introduce $\text{Re}\Sigma_1(\mathbf{k}, \epsilon + i\delta)$, $\text{Re}\Sigma_2(\mathbf{k}, \epsilon + i\delta)$ and $\text{Re}\Sigma_3(\mathbf{k}, \epsilon + i\delta)$ as

$$\text{Re}\Sigma(\mathbf{k}, \epsilon + i\delta) \equiv \text{Re}\Sigma_1(\mathbf{k}, \epsilon + i\delta) + \text{Re}\Sigma_2(\mathbf{k}, \epsilon + i\delta) + \text{Re}\Sigma_3(\mathbf{k}, \epsilon + i\delta) \quad (8)$$

$$\begin{aligned} \text{Re}\Sigma_1(\mathbf{k}, \epsilon + i\delta) &\equiv \int \frac{d^2q}{(2\pi)^2} \int_0^{\infty} \frac{dx}{\pi} [\text{Re}T(\mathbf{q}, x + i\delta) \text{Im}G_0(\mathbf{q} - \mathbf{k}, x - \epsilon - i\delta) \\ &\quad - \text{Im}T(\mathbf{q}, x + i\delta) \text{Re}G_0(\mathbf{q} - \mathbf{k}, x - \epsilon - i\delta)] \end{aligned} \quad (9)$$

$$\text{Re}\Sigma_2(\mathbf{k}, \epsilon + i\delta) \equiv - \int \frac{d^2q}{(2\pi)^2} \int_0^{\epsilon} \frac{dx}{\pi} \text{Re}T(\mathbf{q}, 0) \text{Im}G_0(\mathbf{q} - \mathbf{k}, x - \epsilon - i\delta) \quad (10)$$

$$\text{Re}\Sigma_3(\mathbf{k}, \epsilon + i\delta) \equiv - \int \frac{d^2q}{(2\pi)^2} \int_0^{\epsilon} \frac{dx}{\pi} [\text{Re}T(\mathbf{q}, x + i\delta) - \text{Re}T(\mathbf{q}, 0)] \text{Im}G_0(\mathbf{q} - \mathbf{k}, x - \epsilon - i\delta). \quad (11)$$

It is seen that $\text{Re}\Sigma_1(\mathbf{k}, \epsilon + i\delta)$ is a function of only $\epsilon - vk$ and $\text{Re}\Sigma_2(\mathbf{k}, \epsilon + i\delta)$ a function of only ϵ for the dispersion given by eq. (1). Reflecting the regular behavior of $\text{Re}T(\mathbf{q}, 0)$, $\text{Re}\Sigma_2(\mathbf{k}, \epsilon + i\delta)$ is a regular function of ϵ . Actually for $|\epsilon|, |vk| \ll \epsilon_c$, we obtain

$$\text{Re}\Sigma_1(\mathbf{k}, \epsilon + i\delta) \simeq \text{Re}\Sigma(0, 0) + c_1(\epsilon - vk) \quad (12)$$

$$\text{Re}\Sigma_2(\mathbf{k}, \epsilon + i\delta) \simeq c_2\epsilon, \quad (13)$$

where c_1 and c_2 are constants. On the other hand, we note that $\text{Re}\Sigma_3(\mathbf{k}, \epsilon + i\delta)$ and $\text{Im}\Sigma(\mathbf{k}, \epsilon + i\delta)$ are real and imaginary parts of a function which has a contribution from only the low-energy region, i.e., $|q_x| \ll k_c, |q_y| \ll k'_c, |x| \ll \epsilon_c$.

$$\text{Re}\Sigma_3(\mathbf{k}, \epsilon + i\delta) + i\text{Im}\Sigma(\mathbf{k}, \epsilon + i\delta) \quad (14)$$

$$\simeq - \int_{-\lambda}^{\lambda} \frac{dq_x}{2\pi} \int_{-\lambda'}^{\lambda'} \frac{dq_y}{2\pi} \int_0^{\epsilon} \frac{dx}{\pi} [T(\mathbf{q}, x + i\delta) + U_{\text{eff}}] \text{Im}G_0(\mathbf{q} - \mathbf{k}, x - \epsilon - i\delta), \quad (15)$$

where the cut-offs λ and λ' are small values compared to k_c and k'_c , respectively, and $U_{\text{eff}} \equiv -\lim_{q \rightarrow 0} [\lim_{x \rightarrow 0} T(\mathbf{q}, x + i\delta)]$. We introduce $\Sigma_s(\mathbf{k}, \epsilon + i\delta)$ by subtracting the on-shell value of $\text{Re}\Sigma_3(\mathbf{k}, \epsilon + i\delta)$,

$$\Sigma_s(\mathbf{k}, \epsilon + i\delta) \equiv [\text{Re}\Sigma_3(\mathbf{k}, \epsilon + i\delta) - \text{Re}\Sigma_3(\mathbf{k}, vk + i\delta)] + i\text{Im}\Sigma(\mathbf{k}, \epsilon + i\delta). \quad (16)$$

$\text{Re}\Sigma_s(\mathbf{k}, \epsilon + i\delta)$ is of interest to us, while $\text{Re}\Sigma(\mathbf{k}, \epsilon + i\delta) - \text{Re}\Sigma_s(\mathbf{k}, \epsilon + i\delta) = \text{Re}\Sigma_1(\mathbf{k}, \epsilon + i\delta) + \text{Re}\Sigma_2(\mathbf{k}, \epsilon + i\delta) + \text{Re}\Sigma_3(\mathbf{k}, vk + i\delta)$ is related to various renormalizations such as the shift of the chemical potential, the renormalization of the Fermi velocity, the effects of which are taken into account by replacing v_0 and A_0 in eq. (1) and the renormalization factor Z_0 of the Green function $G_0(\mathbf{k}, \epsilon + i\delta)$. Then, we obtain the renormalized self-energy $\Sigma^*(\mathbf{k}, \epsilon + i\delta)$ and the renormalized Green function $G^*(\mathbf{k}, \epsilon + i\delta)$ as

$$\Sigma^*(\mathbf{k}, \epsilon + i\delta) \equiv [\Sigma_s(\mathbf{k}, \epsilon + i\delta)]_{G_0 \rightarrow G_0^*} \quad (17)$$

$$G^*(\mathbf{k}, \epsilon + i\delta) \equiv \frac{Z_0}{(\epsilon - vk) - \Sigma^*(\mathbf{k}, \epsilon + i\delta)}, \quad (18)$$

where $G_0^*(\mathbf{k}, \epsilon + i\delta) \equiv Z_0/(\epsilon - \xi_k^*)$ and $\xi_k^* \equiv vk_x + Ak_y^2/2$. The subscript $G_0 \rightarrow G_0^*$ on the right hand side (r.h.s.) of eq. (17) indicates that the calculation of $\Sigma_s(\mathbf{k}, \epsilon + i\delta)$ should be performed using the renormalized Green function, $G_0^*(\mathbf{k}, \epsilon + i\delta)$. We note that the on-shell value of $\text{Re}\Sigma_3(\mathbf{k}, \epsilon + i\delta)$, $\text{Re}\Sigma_3(\mathbf{k}, vk + i\delta)$, is of the order of $(vk)^2$, and that this term gives the renormalization of a quasi-particle energy only of the order of k^2 .

Next we need to estimate $\Sigma^*(\mathbf{k}, \epsilon + i\delta)$. Under the conditions of $|q_x| \ll k_c$, $|q_y| \ll k'_c$, $|x| \ll \epsilon_c$, the asymptotic form of $K^*(\mathbf{q}, x + i\delta) - K_0^*$ is given by

$$K^*(\mathbf{q}, x + i\delta) - K_0^* \simeq \begin{cases} \frac{iZ_0^2 x}{4\pi A^{1/2} v(x - vq_x - Aq_y^2/4)^{1/2}} & (x > vq_x + Aq_y^2/4) \\ \frac{Z_0^2 x}{4\pi A^{1/2} v(-x + vq_x + Aq_y^2/4)^{1/2}} & (x < vq_x + Aq_y^2/4), \end{cases} \quad (19)$$

where the particle-particle correlation function $K^*(\mathbf{q}, x + i\delta)$ is calculated using the renormalized Green function $G_0^*(\mathbf{k}, \epsilon + i\delta)$. Using these expressions, we obtain the asymptotic form of $\text{Re}\Sigma^*(\mathbf{k}, \epsilon + i\delta)$ in the limit of $|\epsilon - vk| \ll A^{-1}U_{\text{eff}}^2 Z_0^4 k^2$ for the case of $\epsilon > vk$, for example, as

$$\begin{aligned} & \text{Re}\Sigma^*(\mathbf{k}, \epsilon + i\delta) \\ & \simeq -\frac{Z_0}{4\pi^2 v} \int_{-\lambda'}^{\lambda'} dq_y \int_0^\epsilon dx \text{Re} \left[\frac{-U_{\text{eff}}}{1 + \frac{iU_{\text{eff}} Z_0^2 x}{4\pi A^{1/2} v(Aq_y^2/4 + \epsilon - vk)^{1/2}}} - \frac{-U_{\text{eff}}}{1 + \frac{iU_{\text{eff}} Z_0^2 x}{4\pi A^{1/2} v(Aq_y^2/4)^{1/2}}} \right] \\ & \simeq (\epsilon - vk) \frac{2}{\pi Z_0} \int_0^\infty dq_y \left[\left(\frac{q_y^2}{4} + 1 \right)^{1/2} \arctan \left(\frac{U_{\text{eff}} Z_0^2 \epsilon}{4\pi A^{1/2} v(\epsilon - vk)^{1/2} \left(\frac{q_y^2}{4} + 1 \right)^{1/2}} \right) \right] \end{aligned}$$

$$\begin{aligned}
& - \left(\frac{q_y^2}{4} \right)^{1/2} \arctan \left(\frac{U_{\text{eff}} Z_0^2 \epsilon}{4\pi A^{1/2} v (\epsilon - vk)^{1/2} \left(q_y^2/4 \right)^{1/2}} \right) \Bigg] \\
& \simeq \text{sgn}[k] \frac{(\epsilon - vk)}{Z_0} \int_0^\nu dq_y \left[\left(\frac{q_y^2}{4} + 1 \right)^{1/2} - \left(\frac{q_y^2}{4} \right)^{1/2} \right] \\
& \simeq \text{sgn}[k] \frac{(\epsilon - vk)}{Z_0} \log \nu,
\end{aligned} \tag{20}$$

where

$$\nu \simeq \frac{U_{\text{eff}} Z_0^2 |k|}{A^{1/2} (\epsilon - vk)^{1/2}} \gg 1. \tag{21}$$

The calculations for the case of $\epsilon < vk$ are similar and we obtain the final result in the limit of $|\epsilon - vk| \ll A^{-1} U_{\text{eff}}^2 Z_0^4 k^2$ as

$$\text{Re}\Sigma^*(k, \epsilon + i\delta) \simeq c \frac{(\epsilon - vk)}{Z_0} \log \frac{A|\epsilon - vk|}{U_{\text{eff}}^2 Z_0^4 k^2}, \tag{22}$$

where

$$c = \begin{cases} -\frac{1}{2} & (\epsilon > vk, k > 0) \\ 0 & (\epsilon < vk, k > 0) \\ \frac{1}{2} & (\epsilon > vk, k < 0) \\ 1 & (\epsilon < vk, k < 0) \end{cases}. \tag{23}$$

Reflecting the existence of a logarithmic singularity, the renormalization factor $Z = \lim_{\epsilon \rightarrow vk} (1 - \partial \text{Re}\Sigma / \partial \epsilon)^{-1}$ exhibits an anomalous behavior, $Z = 0$, except in the limit of $\epsilon \rightarrow vk - 0$ in the case of $vk > 0$.

In addition to this, we have evaluated the asymptotic form of $\text{Im}\Sigma^*(k, \epsilon + i\delta)$ in the same limit of $|\epsilon - vk| \ll A^{-1} U_{\text{eff}}^2 Z_0^4 k^2$, which is given by

$$\begin{aligned}
\text{Im}\Sigma^*(k, \epsilon + i\delta) & \simeq \text{Im}\Sigma^*(k, vk + i\delta) \\
& + \frac{(\epsilon - vk)}{Z_0} \left[-\frac{1}{4\pi} \left(\log \frac{A|\epsilon - vk|}{U_{\text{eff}}^2 Z_0^4 k^2} \right)^2 + a \left(\log \frac{A|\epsilon - vk|}{U_{\text{eff}}^2 Z_0^4 k^2} \right) + b \right]
\end{aligned} \tag{24}$$

$$\text{Im}\Sigma^*(k, vk + i\delta) \simeq \frac{U_{\text{eff}}^2 Z_0^4 k^2}{A} \log \frac{U_{\text{eff}} Z_0^2 |k|}{A^{1/2} \epsilon_c^{1/2}} \equiv \Gamma_k, \tag{25}$$

where a and b are constants which are different in each region of $\epsilon > vk$ and $\epsilon < vk$. Especially for $\epsilon \rightarrow vk$, only the first term on the r.h.s. of eq. (24) survives. So far, the real and imaginary parts of the self-energy have been examined, but $\Sigma(k, \epsilon + i\delta)$ must be an analytic function of ϵ . Actually, $\Sigma^*(k, z)$ is given as follows for a complex variable z ($\text{Im } z > 0$) in the limit of $|z - vk| \ll A^{-1} U_{\text{eff}}^2 Z_0^4 k^2$.

$$\begin{aligned}
\Sigma^*(k, z) & \simeq i \text{Im}\Sigma^*(k, vk + i\delta) \\
& + \frac{(z - vk)}{Z_0} \left[-\frac{i}{4\pi} \left(\log \frac{A(z - vk)}{U_{\text{eff}}^2 Z_0^4 k^2} \right)^2 + a' \left(\log \frac{A(z - vk)}{U_{\text{eff}}^2 Z_0^4 k^2} \right) + b' \right],
\end{aligned} \tag{26}$$

where a' and b' are complex constants related to a and b in eq. (24). From the analyticity condition, $\Sigma^*(k, \epsilon + i\delta)$ is interrelated from $\epsilon > vk$ to $\epsilon < vk$ through the upper-half complex plane of ϵ (Fig. 1(c)), and therefore the constants a' and b' in each region are mutually related by a simple relation. When we see $\Sigma^*(k, z)$ as one analytic function of z , the logarithmic term in the real part is related to the second term of the r.h.s. of eq. (24) in the imaginary part. Actually, we obtain $\text{Re } a' = -1/2$ ($1/2$) for $vk > 0$ ($vk < 0$). The leading term of the imaginary part when $\epsilon \simeq vk$, however, is not the second term in eq. (24) but the first one. On the other hand, in the case of $k = k_F$, $\text{Im}\Sigma(k_F, \epsilon + i\delta) \propto \epsilon^2 \log \epsilon$ was obtained,¹⁾ which corresponds to the first term on the r.h.s. of eq. (24), and was the basis of the claim of the Fermi-liquid state. As has been demonstrated, however, the present logarithmic term of the real part cannot be deduced from this imaginary part using the Kramers-Kronig transformation. This has not been noted in previous studies.

Our new result of a logarithmic singular behavior in the real part of the self-energy implies that the renormalization factor Z vanishes in the low-energy limit.

In addition, we note the relationship between the results in the t -matrix approximation and the second-order perturbation theory. When we focus only on the forward scattering process, the self-energy as an analytic function of complex variable z is obtained by the second-order perturbation theory as¹⁰⁾

$$\Sigma^*(k, z) \propto iU^2 z^2 \log \frac{(z - vk)}{\epsilon_c}, \quad (27)$$

and there is no logarithmic term in the real part of the self-energy. In the case of the t -matrix approximation, we found that the self-energy is a logarithmic function not only of $\epsilon - vk$ but also of U_{eff} , as shown in eq. (26). Hence, the logarithmic term in the real part of the self-energy obtained by the t -matrix approximation is a non-perturbative effect of U , which implies that $U = 0$ can be a singular point in two dimensions.

Investigating the derivation in eq. (20) of the logarithmic term in the real part of the self-energy near the on-shell region, we can see that the logarithmic term is related to the property of the t -matrix in the region of $|x - vq_x| < A^{-1}U_{\text{eff}}^2 Z_0^4 k^2$, $Aq_y^2 < A^{-1}U_{\text{eff}}^2 Z_0^4 k^2$, $0 < |x| < |\epsilon|$. This indicates that the on-shell electrons are responsible to the logarithmic term of the self-energy.


Anderson proposed that the special feature of the forward scattering process leads to the vanishing renormalization factor, $Z = 0$, i.e., the breakdown of the Fermi-liquid, and our results have confirmed this based on the conventional many-body perturbation theory using the Feynman diagrams.

So far the calculations of the self-energy have been carried out based on the Green function without damping, Γ_k . The resulting logarithmic behavior in the real part of the self-energy, however, has been seen only in the energy region $|\epsilon - vk| \ll A^{-1}U_{\text{eff}}^2 Z_0^4 k^2 \Gamma_k$, implying the importance of the self-consistent treatment. Actually, Γ_k results in finite Z even when $X \rightarrow 0$, i.e., $G(k, \epsilon + i\delta) \simeq Z_0/[a(X)X + i\Gamma_k]$ with $a(0) \neq 0$. However, $a(X)$ will have a strong dependence on $X = \epsilon - vk$ as


$a(X) = a_0[1 + O(|X|/\Gamma_k)]$, which is different from a conventional Fermi-liquid.

K.Y. acknowledges Hiroshi Kohno for valuable discussions. K.Y. also thanks JSPS Research Fellowships for Young Scientists. This work is financially supported by a Grant-in-Aid for Scientific Research on Priority Area "Anomalous Metallic State near the Mott Transition" (07237102) from the Ministry of Education, Science, Sports and Culture.


-
- [1] C. Hodges, H. Smith and J. W. Wilkins: Phys. Rev. B **4** (1971) 302.
 - [2] P. W. Anderson: Phys. Rev. Lett. **64** (1990) 1839; *ibid.* **65** (1990) 2306.
 - [3] P. W. Anderson: Phys. Rev. Lett. **71** (1993) 1220.
 - [4] M. Fabrizio, A. Parola and E. Tosatti: Phys. Rev. B **44** (1991) 1033.
 - [5] P. C. E. Stamp: J. Phys. I (France) **3** (1993) 625
 - [6] J. R. Engelbrecht and M. Randeria: Phys. Rev. Lett. **65** (1990) 1032; Phys. Rev. B **45** (1992) 12419.
 - [7] H. Fukuyama and Y. Hasegawa: Prog. Theor. Phys. Suppl. **101** (1990) 441.
 - [8] H. Fukuyama, Y. Hasegawa and O. Narikiyo: J. Phys. Soc. Jpn. **60** (1991) 372; *ibid.* 2013; O. Narikiyo: Thesis, University of Tokyo, 1992.
 - [9] O. Narikiyo and K. Miyake: J. Phys. Soc. Jpn. **64** (1995) 2290.
 - [10] H. Fukuyama and M. Ogata: J. Phys. Soc. Jpn. **63** (1994) 3923.



(a)



(b)



(c)